

B

Name:

Math Models of Operations Research, MATP 4700/ DSES 4770.

First Exam, Tuesday, September 30, 2008.

You may use any result from your notes or a homework that is clearly stated. You may use one sheet of handwritten notes, but no other sources. The exam consists of five questions, and lasts one hundred and ten minutes.

Q1	/ 15
Q2	/ 21
Q3	/ 20
Q4	/ 24
Q5	/ 20
Total	/ 100
Grade	

SOLUTIONS.

1. (15 points; each part is worth 5 points.)

The following tableau represents a linear program in standard form:

	↓	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$
3	0	0	0	0	2	0	
10	-3	0	0	1	10	0	
12/8	12	8	0	-2	0	6	1
1/1	1	1	1	-1	0	0	0

- (a) Give an optimal basic feasible solution to this linear program.
- (b) Find another optimal basic feasible solution.
- (c) Show that the set of optimal solutions is unbounded.

(a)  $x = (0, 1, 0, 10, 0, 12)$

(b) Bring  $x_1$  into basis. New tableau:

3	0	0	0	0	2	0
13	0	3	-3	1	10	0
4	0	-4	2	0	6	1
1	1	1	-1	0	0	0

Alternate optimal b.f.s:

$x = (1, 0, 0, 13, 0, 4)$ .

(c)  $x_3$  has reduced cost of zero.  
 In original tableau, if try to introduce  $x_3$  into basis, find a ray.  
 So set of optimal solutions includes

$$x = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 10 \\ 0 \\ 12 \end{bmatrix} + t \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 2 \end{bmatrix} \text{ for any } t \geq 0.$$

2. (21 points. Each part is worth 7 points.)

The following tableau represents a linear program in standard form:

	$x_1$	$x_2$	$x_3$	$x_4$
3	-3	0	-1	0
0	4	1	-1	0
12	12	0	3	1

- (a) There are two nonbasic variables. What are the simplex directions corresponding to each of these nonbasic variables?
- (b) Find the steplengths in the directions found in part 2a. Hence find the new basic feasible solutions that would be obtained by introducing each of these nonbasic variables.
- (c) Show that the optimal solution can be found in one iteration by choosing the entering variable appropriately.

(a) Introduce  $x_1$ :  $d^{(1)} = \begin{bmatrix} 1 \\ -4 \\ 0 \\ -12 \end{bmatrix}$       Introduce  $x_2$ :  $d^{(2)} = \begin{bmatrix} 0 \\ 1 \\ 1 \\ -3 \end{bmatrix}$

(b) Min ratios give 0 for  $d^{(1)}$ ,  $\frac{1}{3}$  for  $d^{(2)}$

(c) Introduce  $x_3$ :

7	1	0	0	$\frac{1}{3}$
4	8	1	0	$\frac{1}{3}$
4	4	0	1	$\frac{1}{3}$

Optimal soln is  $x = (0, 4, 4, 0)$ .

3. (20 points)

Consider the following tableau for a standard form linear program:

$$\Pi = \begin{array}{c|cccccccc} & x_1 & x_2 & x_3 & x_4 & x_5 & x_6 & x_7 & x_8 \\ \hline 3 & -3 & -2 & -1 & 0 & -2 & 0 & -4 & -6 \\ \hline 3 & 7 & 13 & -2 & 1 & 6 & 0 & 2 & 3 \\ \hline 1 & -3 & -6 & 1 & 0 & 10 & 1 & 3 & -1 \end{array}$$

Several simplex pivots are taken. The pivot matrix corresponding to this sequence of pivots is:

$$P = \begin{bmatrix} 1 & 6 & 13 \\ 0 & 1 & 2 \\ 0 & 3 & 7 \end{bmatrix}$$

The new basic sequence consists of  $x_1$  and  $x_3$ .

- (a) (6 points) What is the basic feasible solution obtained as a result of this sequence of pivots? What is its value?
- (b) (7 points) Show that the solution you obtained in part 3a is not optimal.
- (c) (7 points) How would you next update the pivot matrix?

↓

$$\text{PM} = \begin{array}{c|cccccccc} 34 & 0 & -2 & 0 & 6 & 164 & 13 & 47 & -1 \\ \hline 5 & & 1 & & & & & & \\ \hline 16 & & -3 & & & & & & \end{array}$$

(a)  $x_1 = 5, x_3 = 16, \text{ other } x_i = 0. \text{ Value} = -34$

(b)  $x_2$  and  $x_8$  have negative reduced cost, so not optimal.

(c) We only examine  $x_2$  entering basis.

$x_1$  leaves basis.

New  $P = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & 0 \\ 0 & 3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 6 & 13 \\ 0 & 1 & 2 \\ 0 & 3 & 7 \end{bmatrix}$

4. (24 points; each part is worth 6 points)

The variables in the following linear programming problem have upper bounds:

$$\begin{array}{llll}
 \min & 5x_1 & - & 3x_2 \\
 \text{subject to} & 3x_1 & - & 2x_2 + x_3 = 16 \\
 & 2x_1 & + & x_2 + x_4 = 9 \\
 & & & 0 \leq x_1 \leq 4 \\
 & & & 0 \leq x_2 \leq 3 \\
 & & & 0 \leq x_3 \leq 25 \\
 & & & 0 \leq x_4 \leq 4
 \end{array}$$

Slack variables are not introduced for the upper bound constraints. Instead, variables can be nonbasic at either their upper or lower bounds.

- (a) Show that  $x_3$  must be basic in any basic feasible solution. (Hint: Consider the bounds on  $x_1$  and  $x_2$ .)
- (b) Find all the basic feasible solutions where at least one variable is at its upper bound.
- (c) Show that the original linear program is equivalent to the following linear program:

$$\begin{array}{ll}
 \min & 5x_1 - 3x_2 \\
 \text{subject to} & 5 \leq 2x_1 + x_2 \leq 9 \\
 & 0 \leq x_1 \leq 4 \\
 & 0 \leq x_2 \leq 3
 \end{array}$$

(For the purposes of this question, two linear programs are equivalent if any feasible solution in one can be mapped to a feasible solution in the other with the same objective value.)

- (d) Plot the feasible region of the linear program in part 4c. Mark each of the basic feasible solutions you found in part 4b. Find the optimal solution graphically.

(a)  $3x_1 - 2x_2 \leq 12 \Rightarrow x_3 \geq 4$   
 $3x_1 - 2x_2 \geq -6 \Rightarrow x_3 \leq 22$  } So  $x_3$  lies strictly between its bounds, so always basic


(b)

$x_1$	$x_2$	$x_3$	$x_4$	Point	Status
B	LB	B	UB	$x = (2\frac{1}{2}, 0, 8\frac{1}{2}, 4)$	BFS
B	UB	B	LB	$x = (3, 3, 13, 0)$	BFS
B	UB	B	UB	$x = (1, 3, 19, 4)$	BFS
LB	B	B	UB	$x = (0, 5, 16, 4)$	violates $x_2$ bound
UB	B	B	LB	$x = (4, 1, 6, 0)$	BFS
UB	B	B	UB	$x = (4, -3, , 4)$	violates $x_2$ bound
LB	UB	B	B	$x = (0, 3, , 6)$	violates $x_4$ bound
UB	LB	B	B	$x = (4, 0, 4, 1)$	BFS
UB	UB	B	B	$x = (4, 3, , -2)$	violates $x_4$ bound.

} So 5 BFSs.

(intentionally left blank)

(c) First constraint is redundant, since bounds on  $x_3$  always bind.  
 Second constraint and bounds on  $x_4$  give constraint  
 in second formulation.



Optimal solution:  $x = (1, 3, 19, 4)$ .

5. (20 points; each part is worth 5 points)

A company manufactures parts  $i = 1, \dots, m$  in weeks  $t = 1, \dots, n$ , where each unit of part  $i$  requires  $a_{i,k}$  units of production resource  $k = 1, \dots, q$  and has value  $v_i$ . Production resource capacities  $b_k$  cannot be exceeded in any period and part demands  $d_{i,t}$  must be met. Express each of the following as linear constraint(s) in these parameters and the nonnegative decision variables  $x_{i,t} :=$  number of units of  $i$  produced in week  $t$  and  $z_{i,t} :=$  inventory of product  $i$  held at the end of period  $t$ . Initial inventories all = 0.

- (a) No production capacity can ever be exceeded.
- (b) The total value of held inventories should never exceed 200.
- (c) Quantities of each part  $i$  available after week 1 should balance with demand and accumulated inventory.
- (d) Quantities of each part  $i$  available after weeks  $2, \dots, n - 1$  should balance with demand and accumulated inventory.

See Version A.